

Innovations in Wound Care

# The role of wound cleansing in the management of wounds

Brought to you by



Sponsored by



# Faculty

*Speaker*

**Martha Kelso, RN, HBOT, CEO, WCP**  
Wound Care Plus, LLC



*Moderator*

**Melissa Warner, EVP**  
Wound Care Advisor



# Agenda For Today

This 30-minute presentation will feature learning opportunities that will provide in-depth instruction and demonstration in wound care treatments. After this webinar, the learner will be able to:

- Identify the role of proper wound cleansing
- Discuss how to select and use non-toxic wound cleansers
- Describe advantages of collagen for managing a chronic wound

# Objectives

At the end of this webinar, the learner will be able to:

- **Identify the role of proper wound cleansing**
- **Discuss how to select and use non-toxic wound cleansers**

**A) The wound microenvironment**

**B) Is your wound cleansing practice up to date?**

- Cleansing and its role
- Basic Cleansing Techniques
- When not to clean a wound

**C) The ideal wound cleanser**

**D) The use of hypochlorous acid as a wound cleanser**

**E) Clinical Case studies**



# **THE WOUND MICROENVIRONMENT**

# Wound microenvironment of chronic wounds represents a major therapeutic challenge<sup>1</sup>

The most relevant factors that influence the **healing process** are:




(1) Kruse CR et al (2015): *Wound Repair and Regeneration* 23(4): 456–464 - (2) Caley MP et al (2015): *Advances in Wound Care* 4(4): 225-234

(3) Watters C et al (2015): *Chronic Wound Care Management and Research* 2: 53–62 - (4) Castilla DM et al (2012): *Advances in Wound Care* 1(6): 225-230

# The importance of pH in wound healing




Wounds with an alkaline pH have demonstrated lower rates of healing<sup>6</sup>



(5) Lambers H et al (2006): *J Cosmet Sci* 2006; 28: 359–370 – (6) Gethin G (2007): *Wounds UK*, 2007:3/3

# The importance of acidic pH

“An **acidic environment** in a wound bed is an additional benefit that can contribute to **reboot the wound** towards healing”







**IS YOUR WOUND CLEANSING  
PRACTICE UP TO DATE?**

# Is your wound cleansing practice up to date?

## “Cleansing and its role”

Wound cleansing is performed to remove surface contaminants, **bacteria, non-viable tissue** and excess **exudate** from the wound bed and surrounding skin <sup>10</sup>



**Wound cleansing** can be an effective way to **remove inflammatory stimulants** and **local barriers** from the **wound bed**<sup>10</sup>

An **ideal wound cleanser** should **modulate the wound microenvironment** balancing the management of key components with **preservation of tissue safety**<sup>11</sup>

# Is your wound cleansing practice up to date?


## “Basing cleansing techniques”

### Swabbing

(use items that don't leave debris in wound bed)



### Irrigation



# Is your wound cleansing practice up to date?

## “When not to clean a wound ”

Dry gangrenous wounds  
(want to keep dry)



Fragile granulation tissue  
(may be traumatic)





# **THE IDEAL WOUND CLEANSER**

## The ideal wound cleanser

When you ask your patient about their basic wound care at home.... what is the usual (and unfortunate) answer?



# The ideal wound cleanser

## “Antiseptic”

- Research has shown that **antiseptics** have a **negative impact on healing wounds** (Atiyeh, et.al. Int Wound J. 2009)
- Antiseptic categories include **alcohols, iodine, Chlorhexidine Gluconate (CHG), silver, hydrogen peroxide**
- Antiseptics can be used in the right situations, and durations



# The ideal wound cleanser

There are several broad categories of solutions that can be used:

Wound cleansing can help to achieve the goals of **wound bed preparation** by removing microorganism, biological and environmental debris<sup>12</sup>

## HYPOCHLOROUS ACID

**Hypochlorous acid** is produced by the body's immune cells in response to invading pathogens. When used as wound cleanser ingredient, it acts as a preservative by **inhibiting the growth of microorganisms** within the solution<sup>12</sup>

## SALINE SOLUTION

**Saline solution does not generally contain a preservative**, so bacterial growth can occur once exposed to opportunistic microorganisms<sup>12</sup>

## SODIUM HYPOCHLORITE

**Sodium hypochlorite** is familiar to HCPs as Dakin's solution (0.5% sodium hypochlorite). Dakin's solution can be **injurious to the wound tissue** and **can slow down wound healing**<sup>12</sup>



# The ideal wound cleanser

The characteristics of an ideal wound cleanser



- The ideal wound cleanser should be **non-cytotoxic** to tissue

---

- The ideal wound cleanser should **decrease colonization of the wound bed**

---

- The ideal wound cleanser should be **cost-effective** and **stable**

---

- The ideal wound cleanser should **not be an alkaline pH**




# **THE USE OF HYPOCHLOROUS ACID AS A WOUND CLEANSER**

# The use of hypochlorous acid as a wound cleanser

Hypochlorous acid is one of the major inorganic bactericidal compound of **innate immunity** and it is **effective against a broad range of microorganisms**

Although they look similar, **hypochlorous acid** and **sodium hypochlorite** are still **very different**



The cell wall of pathogens is **negatively charged**



**HClO can easily penetrate the cell wall and destroy the pathogens from the inner of the cell.**

**ClO<sup>-</sup>/sodium hypochlorite can not easily penetrate the wall of pathogens**



At the same concentration, the biocidal activity of **HClO is 80 times stronger than ClO<sup>-</sup>**

# The use of hypochlorous acid as a wound cleanser

	<b>HYPOCHLOROUS ACID HClO</b>	<b>SODIUM HYPOCHLORITE NaClO</b>
<b>SAFETY</b>	<ul style="list-style-type: none"><li>✓ HClO is <b>naturally</b> found in our body<sup>13</sup></li><li>✓ It is produced by the <b>immune cells</b> (white blood cells) in response to invading pathogens<sup>13</sup></li></ul>	<ul style="list-style-type: none"><li>✗ NaClO can be <b>injurious to the wound</b><sup>15</sup></li><li>✗ NaClO can be <b>cytotoxic</b><sup>15</sup></li><li>✗ NaClO can <b>slow down wound healing</b><sup>15</sup></li></ul>
<b>EFFICACY</b>	<ul style="list-style-type: none"><li>✓ HClO is <b>80 times stronger</b> than NaClO<sup>14</sup></li></ul>	<ul style="list-style-type: none"><li>✗ NaClO <b>does not easily penetrates</b> the bacteria<sup>14</sup></li></ul>
<b>pH</b>	<ul style="list-style-type: none"><li>✓ <b>Acidic pH</b></li></ul>	<ul style="list-style-type: none"><li>✗ <b>Alkaline pH</b></li></ul>



# **CLINICAL CASE STUDIES: LEG ULCERS**

# Clinical cases (1/3)

- AGE, SEX: 48, male
- WOUND ONSET: 15 years before treatment with a **wound cleanser containing hypochlorous acid**
- COMORBIDITY: severe
  - Essential hypertension
  - Lymphoma: in 1999
  - Epilepsy
  - Valvular heart disease
  - Osteomyelitis of the knee in 2010
- CASE HISTORY BEFORE NEXODYN:
  - Traffic accident with left lower limb severe injury. Subsequent alteration of vascular architecture with the appearance of a large ulcer on the lower third of the left leg after chemotherapy, due to a car accident about 15 years ago
  - Large ulcer on the left leg (lower 1/3) treated with cycles of hyperbaric oxygen therapy. Since then, the lesion has never come to resolution.
  - From November 2011 to June 2013, the patient underwent 4 surgeries with engineered graft and skin graft without any benefit, with the exception of wound depth, becoming superficial.

# Clinical cases (1/3)

Large ulcer on the left leg (lower 1/3)



From T0  
to ≈ 10 months



Time (days)	Tissues	Exudate	Depth	Area (cm <sup>2</sup> )	VAS (pain)
0	Colonized	Hyperexudating	Superficial	250	7
12	Colonized	Average	Superficial	250	6
42 (≈1.5 m)	Cleansed	Controlled	Superficial	215	4
183 (≈6 m)	Re-epithelising	Controlled	Superficial	184	2
302 (≈10 m)	Re-epithelising	Controlled	Superficial	135	2
<b>Result</b>	<b>Improved</b>	<b>Improved</b>	<b>Unchanged</b>	<b>-46.0%</b>	<b>-71.4%</b>

# Clinical cases (2/3)

- AGE: 31
- SEX: male
- WOUND ONSET: 2 years before treatment with a **wound cleanser containing hypochlorous acid**
- COMORBIDITY: severe
  - Young refractory severe obesity; sleeve gastrectomy in 2010 (pre-surgery weight 227 kg; 148 kg in 2011)
  - Psoriatic arthritis
  - Very large peripheral and lower edemas on a lymphostatic basis
- CASE HISTORY:
  - Stasis ulcer that does not tolerate the bandage
  - DE grafting with improvement on 06/2013
  - Very low compliance
  - Constant infections



# Clinical cases (2/3)

Stasis ulcer that does not tolerate the bandage



From T0  
to ≈ 9.5 months after



Time (days)	Tissues	Exudate	Depth	Area	VAS (pain)
0	Infected	Hyperexudating	Deep	300	8
22	Cleansed	Average	Superficial	250	5
155 (≈5 m)	Cleansed	Average	Superficial	180	2
236 (≈7.5 m)	Cleansed	Hyperexudating	Superficial	150	2
295 (≈9.5 m)	Cleansed	Poor	Superficial	118	2
<b>Result</b>	<b>Improved</b>	<b>Improved</b>	<b>Improved</b>	<b>-60.66%</b>	<b>-75%</b>

# Clinical cases (3/3)

- AGE: 83
- SEX: male
- WOUND ONSET: 2.5 years before treatment with a **wound cleanser containing hypochlorous acid**
- COMORBIDITY: severe
  - Benign prostatic hyperplasia with previous TURP (Transurethral resection of the prostate)
  - Hypertension
  - Post-thrombotic syndrome on the right leg
  - Lipotimic episodes
  - Paroxysmal atrial fibrillation
  - Chronic carential anaemia
- CASE HISTORY:
  - April 2012: hospitalization with a diagnosis of bilateral ulcers of the lower limbs by pyoderma gangrenosus and polymicrobial super-infection, with severe sepsis, eurhythmic paroxysmal atrial fibrillation by wandering pacemaker at heparin in coagulant dosage, mild heart failure, anasarctic condition in severe protein-caloric malnutrition, psychomotor agitation with delirium perhaps iatrogenic in nature (carbapenem and opiates), suspected sleep apnea syndrome.
  - April 2013: new hospitalization for re-grafting

# Clinical cases (3/3)

Bilateral ulcers of the lower limbs



Man, 83 yo

From T0  
to ≈ 9 months after



Time (days)	Tissues	Exudate	Depth	Area (cm <sup>2</sup> )	VAS (pain)
-455 (≈15 m)	Slough	Hyperexudating	Superficial	160	6
-70 (≈2 m)	Slough	Hyperexudating	Superficial	140	6
0	Slough	Hyperexudating	Superficial	120	5
<b>Result</b>	<b>Unchanged</b>	<b>Unchanged</b>	<b>Unchanged</b>	<b>-25%</b>	<b>-17%</b>
<b>START OF TREATMENT WITH NEXODYN</b>					
0	Slough	Hyperexudating	Superficial	120	5
190 (≈6 m)	Slough	Average	Superficial	70	2
275 (≈9 m)	Slough	Poor	Superficial	38	1
<b>Result</b>	<b>Improved</b>	<b>Improved</b>	<b>Unchanged</b>	<b>-68.33%</b>	<b>-80%</b>

**Thank you to Angelini  
for sponsoring this webinar**



**ANGELINI**

We will take a few minutes to review.....

# Product features

NEXODYN® can support the physiological healing process

NEXODYN® is a FDA-cleared **hypochlorous acid-based wound cleanser**, developed for topical treatment in the field of **acute** and **chronic wound management**

## MAIN PRODUCT FEATURES

- ✓ **Acidic pH (2.5 – 3.0)**
- ✓ **High purity (>95% of free chlorine species derived from HClO)**
- ✓ **Free Chlorine species: 40-70 ppm**
- ✓ **Long stability: 30 days from first opening**



The mechanical action of the fluid flowing across the lesion can help to **remove biologic and inert materials** such as **microorganisms, biological debris** and **environmental dirt**

# Bacterial activity tests

The antimicrobial preservative effectiveness of HClO has been demonstrated against the organisms in the table below in *in vitro* testing (Time Kill Assay):

Pathogenic Bacteria	Log Reduction / Exposure Time
<i>Staphylococcus aureus</i>	99.9992% (5.11 Log <sub>10</sub> ) after 15 sec
<i>Staphylococcus pyogenes</i>	99.9958% (4.38 Log <sub>10</sub> ) after 15 sec
<i>Staphylococcus epidermidis</i>	99.9499% (3.30 Log <sub>10</sub> ) after 15 sec
<i>Pseudomonas aeruginosa</i>	>99.9999% (> 6.11 Log <sub>10</sub> ) after 15 sec
<i>Escherichia coli</i>	>99.999% (> 5.55 Log <sub>10</sub> ) after 15 sec
<i>Multi-drug resistant (MDR) Staphylococcus aureus</i>	>99.999% (> 5.44 Log <sub>10</sub> ) after 15 sec
<i>Extended-spectrum beta-lactamase (ESBL) producing Enterobacteriaceae</i>	>99.9999% (> 6.23 Log <sub>10</sub> ) after 15 sec
<i>Vancomycin intermediate resistant Staphylococcus aureus (VISA)</i>	>99.999% (>5.84 Log <sub>10</sub> ) after 15 sec
<i>Multi-drug resistant (MDR) and OXA-48 producing Klebsiella pneumoniae</i>	>99.999% (> 5.32 Log <sub>10</sub> ) after 15 sec
<i>Extended-spectrum beta-lactamase (ESBL) producing Proteus mirabilis</i>	>99.999% (>5.99 Log <sub>10</sub> ) after 15 sec
<i>Multi-drug resistant (MDR) Escherichia coli</i>	>99.999% (>5.92 Log <sub>10</sub> ) after 15 sec
<i>Candida albicans</i>	>99.999% (>5.01 Log <sub>10</sub> ) after 15 sec

# How to use NEXODYN®

Applying NEXODYN® on wounds is **fast and simple**



**1** At each medication, the whole lesion area should be abundantly sprayed with Nexodyn™



**2** The solution should be allowed to dry. No rinsing required.



**3** A second application of Nexodyn™ can be consecutively repeated, if necessary



**4** The solution should be allowed to dry. No rinsing required.



**5** Following cleansing with Nexodyn™ standard therapy can be applied as required.

Contraindications: Do not use in case of hypersensitivity to any component of the product (hypochlorous acid, chlorine and hypochlorite ion)

# The right tools make all the difference for your patients!








Wholesale Medical Supplies  
NATIONWIDE DISTRIBUTION

☎ 1-800-461-1370

Delivered to your patient's home



IMPACT MEDICAL SUPPLY



WHERE TO BUY

# References

- (1) Kruse CR et al (2015): *Wound Repair and Regeneration* 23(4): 456–464
- (2) Caley MP et al (2015): *Advances in Wound Care* 4(4): 225-234
- (3) Watters C et al (2015): *Chronic Wound Care Management and Research* 2: 53–62
- (4) Castilla DM et al (2012): *Advances in Wound Care* 1(6): 225-230
- (5) Lambers H et al (2006): *J Cosmet Sci* 2006; 28: 359–370
- (6) Gethin G (2007): *Wounds UK*, 2007:3/3
- (7) Greener B et al (2005): *J Wound Care* 14(2): 59–61
- (8) Hostacka´ A et al (2010): *Folia Microbiol.* 55 (1): 75–78
- (9) Basavraj S et al (2015): *Wounds* 27(1):5-11
- (10) McLain N and Moore Z (2015): *Cochrane Database of Systematic Review*, 4 (CD01167)
- (11) Main RC (2008): *J Wound Care* 17(3): 112-114
- (12) Wolcott R et al (2014): *Wounds International* (3): 25-31
- (13) Armstrong DC et al (2015): *Ostomy Wound Manage* 61(5):S2-S19
- (14) Rossi-Fedele G (2011): *J Int. Endod. J* 44: 792-799
- (15) Hildago E et al (2002): *Chemico-Biological Interactions* 139: 265-282
- (16) Kelso, Martha. (March 16, 2018). The Role of Hypochlorous Acid in Managing Wounds: Reduction in Antibiotic Usage. Retrieved from <http://www.woundsource.com/blog/role-hypochlorous-acid-in-managing-woundsreduction-in-antibiotic-usage>



QUESTION!